

An analytical exercise*

Leonhard Euler[†]

1. In considering the infinite product for expressing cosine in terms of any angle, which is

$$\cos \frac{\pi}{2n} = \left(1 - \frac{1}{nn}\right) \left(1 - \frac{1}{9nn}\right) \left(1 - \frac{1}{25nn}\right) \left(1 - \frac{1}{49nn}\right) \text{ etc.},$$

it comes to mind to investigate a method which itself works by the nature of the value of this product, which we in fact already know to be equal to $\cos \frac{\pi}{2n}$. Indeed, it is possible, though with many labors which were brought about by artifice, for a geometric demonstration of this to be given which I do not consider at all unpleasant.

2. Therefore I set,

$$S = \left(1 - \frac{1}{nn}\right) \left(1 - \frac{1}{9nn}\right) \left(1 - \frac{1}{25nn}\right) \text{ etc.},$$

and then the logarithms produced by me are summed:

$$lS = l\left(1 - \frac{1}{nn}\right) + l\left(1 - \frac{1}{9nn}\right) + l\left(1 - \frac{1}{25nn}\right) + \text{ etc.},$$

*Delivered October 3, 1776. Originally published as *Exercitatio analytica*, Nova Acta Academiae Scientiarum Imperialis Petropolitinae **8** (1794), 69-72, and republished in *Leonhard Euler, Opera Omnia*, Series 1: Opera mathematica, Volume 16, Birkhäuser, 1992. A copy of the original text is available electronically at the Euler Archive, at www.eulerarchive.org. This paper is E664 in the Eneström index.

[†]Date of translation: December 8, 2004. Translated from the Latin by Jordan Bell, 2nd year undergraduate in Honours Mathematics, School of Mathematics and Statistics, Carleton University, Ottawa, Ontario, Canada. Email: jbell3@connect.carleton.ca. Part of this translation was written during an NSERC USRA supervised by Dr. B. Stevens.

and since

$$l(1 - \frac{1}{x}) = -\frac{1}{x} - \frac{1}{2xx} - \frac{1}{3x^3} - \frac{1}{4x^4} - \text{etc.},$$

the successive series set out with their signs changed are:

$$\begin{aligned} -lS &= \frac{1}{nn} + \frac{1}{2n^4} + \frac{1}{3n^6} + \frac{1}{4n^8} + \text{etc.} \\ &+ \frac{1}{9nn} + \frac{1}{2 \cdot 9^2 n^4} + \frac{1}{3 \cdot 9^3 n^6} + \frac{1}{4 \cdot 9^4 n^8} + \text{etc.} \\ &+ \frac{1}{25nn} + \frac{1}{2 \cdot 25^2 n^4} + \frac{1}{3 \cdot 25^3 n^6} + \frac{1}{4 \cdot 25^4 n^8} + \text{etc.} \\ &+ \frac{1}{49nn} + \frac{1}{2 \cdot 49^2 n^4} + \frac{1}{3 \cdot 49^3 n^6} + \frac{1}{4 \cdot 49^4 n^8} + \text{etc.} \\ &\text{etc.} \end{aligned}$$

3. But now if we arrange each of the rows vertically in columns, we will obtain the following series for $-lS$:

$$\begin{aligned} -lS &= \frac{1}{nn} (1 + \frac{1}{3^2} + \frac{1}{5^2} + \frac{1}{7^2} + \frac{1}{9^2} + \text{etc.}) \\ &+ \frac{1}{2n^4} (1 + \frac{1}{3^4} + \frac{1}{5^4} + \frac{1}{7^4} + \frac{1}{9^4} + \text{etc.}) \\ &+ \frac{1}{3n^6} (1 + \frac{1}{3^6} + \frac{1}{5^6} + \frac{1}{7^6} + \frac{1}{9^6} + \text{etc.}) \\ &+ \frac{1}{4n^8} (1 + \frac{1}{3^8} + \frac{1}{5^8} + \frac{1}{7^8} + \frac{1}{9^8} + \text{etc.}) \\ &\text{etc.} \end{aligned}$$

Thus in this way, the investigation is led to the summation of series of even powers of the harmonic progression, $1, \frac{1}{3}, \frac{1}{5}, \frac{1}{7}, \text{etc.}$

4. By putting $\frac{\pi}{2} = \rho$ for the sake of brevity, it can then be seen that the sums of powers can be represented in the following manner:

$$\begin{aligned} 1 + \frac{1}{3^2} + \frac{1}{5^2} + \frac{1}{7^2} + \text{etc.} &= A\rho^2, \\ 1 + \frac{1}{3^4} + \frac{1}{5^4} + \frac{1}{7^4} + \text{etc.} &= B\rho^4, \\ 1 + \frac{1}{3^6} + \frac{1}{5^6} + \frac{1}{7^6} + \text{etc.} &= C\rho^6, \\ &\text{etc.} \end{aligned}$$

with the first one taken as $A = \frac{1}{2}$, and then the other letters determined by the preceding ones in the following way:

$$B = \frac{2}{3}A^2, C = \frac{2}{5} \cdot 2AB, D = \frac{2}{7}(2AC + BB),$$

$$E = \frac{2}{9}(2AD + 2BC), F = \frac{2}{11}(2AE + 2BD + CC), \text{etc.}$$

the truth of which will soon shine forth from a most beautiful conjunction of analysis.

5. With the substituted values, this series is obtained:

$$-lS = \frac{a\rho^2}{nn} + \frac{1}{2} \cdot \frac{B\rho^4}{n^4} + \frac{1}{3} \cdot \frac{C\rho^6}{n^6} + \frac{1}{4} \cdot \frac{D\rho^8}{n^8} + \text{etc.}$$

If we put $\frac{\rho}{n} = x$, so that $x = \frac{\pi}{2n}$, then this series is obtained:

$$-lS = Axx + \frac{1}{2}Bx^4 + \frac{1}{3}Cx^6 + \frac{1}{4}Dx^8 + \text{etc.}$$

So that we can get rid of the fractions $\frac{1}{2}, \frac{1}{3}, \frac{1}{4}$, etc., we differentiate, and then after dividing the result by $2\partial x$ it follows,

$$-\frac{\partial S}{2S\partial x} = Ax + Bx^3 + Cx^5 + Dx^7 + \text{etc.}$$

6. Then for the sake of brevity we set $-\frac{\partial S}{2S\partial x} = t$, so that we have

$$t = Ax + Bx^3 + Cx^5 + Dx^7 + \text{etc.},$$

for which the square is seen to be this series:

$$\begin{array}{llll} tt = A^2xx & + 2ABx^4 + 2ACx^6 & + 2ADx^8 + 2AEx^{10} & + \text{etc.} \\ & + BB & + 2BC + 2BD & + \text{etc.} \\ & & + CC & + \text{etc.} \end{array}$$

and in this, each of the powers of x are the very same as those found in the formulas which express the determination of the letters A, B, C, D : the only thing missing are the coefficients $\frac{2}{3}, \frac{2}{5}, \frac{2}{7}$, etc.

7. But we can indeed introduce these coefficients by integrating after we have multiplied through by $2\partial x$. So it will be obtained that

$$2 \int tt\partial x = \frac{2}{3}A^2x^3 + \frac{2}{3} \cdot 2ABx^5 + \frac{2}{7}(2AC + BB)x^7 + \frac{2}{9}(2AD + 2BC)x^9 + \frac{9}{11}(2AE + 2BD + CC)x^{11} + \text{etc.}$$

Since at present,

$$\frac{2}{3}A^2 = B, \frac{2}{3} \cdot 2AB = C, \frac{2}{7}(2AC + BB) = D, \text{etc.},$$

with these values substituted in, we then attain this series:

$$2 \int tt\partial x = Bx^3 + Cx^5 + Dx^7 + Ex^9 + \text{etc.}$$

8. Therefore with the series we had from before:

$$t = Ax + Bx^3 + Cx^5 + Dx^7 + \text{etc.},$$

this equation clearly follows:

$$t = Ax + 2 \int tt\partial x,$$

which when differentiated yields

$$\partial t = A\partial x + 2tt\partial x = \frac{1}{2}\partial x + 2tt\partial x, \text{ because } A = \frac{1}{2}.$$

From this we therefore have $2\partial t = \partial x(1 + 4tt)$, from which it is $\partial x = \frac{2\partial t}{1+4tt}$, whose integral is at hand, namely $x = A \text{ tang.} 2t$,¹ for which the addition of a constant is not desired, since by setting $x = 0$ then t should simultaneously vanish. Hence with this equation that has been discovered, if the quantity x is seen as an angle then it will be $2t = \text{tang.} x$. Since it was already $t = -\frac{\partial S}{2S\partial x}$, from this are deduced these equations:

$$-\frac{\partial S}{S\partial x} = \text{tang.} x, \text{ and } -\frac{\partial S}{S} = \frac{\partial x \sin x}{\cos x}.$$

¹Translator: Euler uses here A tang. $2t$ to express the arctangent of $2t$, and later uses tang. x to stand for the tangent of x .

9. Therefore, since it is $\partial x \sin x = -\partial \cos x$, it will be $\frac{\partial S}{S} = \frac{\partial \cos x}{\cos x}$, and then by integrating, $lS = l \cos x + C$, which to be consistent should be defined such that setting $x = 0$ makes $lS = 0$. Thus it will be $C = 0$, so $lS = l \cos x$, and then so that the numbers can come forth, $S = \cos x$.

10. Moreover, since we already had that $x = \frac{\pi}{2n}$, from this it is clear that the value being sought out for S is revealed to be $S = \cos \frac{\pi}{2n}$, which agrees totally with what was known from before. Therefore, analysis most excellently confirms the former relation between the letters A, B, C, D , which I introduced in earlier calculations.